

兔抗 EIF3D 多克隆抗体

中文名称：兔抗 EIF3D 多克隆抗体

英文名称：Anti-EIF3D rabbit polyclonal antibody

别 名：EIF3S7; eIF3-p66; eIF3-zeta

抗 原：EIF3D

储 存：冷冻 (-20°C) 避光

宿 主：Rabbit

相关类别：一抗

反应种属：Human Mouse

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	Eukaryotic translation initiation factor-3 (eIF3), the largest of the eIFs, is a multiprotein complex composed of at least ten nonidentical subunits. The complex binds to the 40S ribosome and helps maintain the 40S and 60S ribosomal subunits in a dissociated state. It is also thought to play a role in the formation of the 40S initiation complex by interacting with the ternary complex of eIF2/GTP/methionyl-tRNA, and by promoting mRNA binding.
Applications:	WB, IHC
Name of antibody:	EIF3D
Immunogen:	Synthesized peptide derived from internal of human EIF3D.
Full name:	eukaryotic translation initiation factor 3, subunit D
Synonyms :	EIF3S7; eIF3-p66; eIF3-zeta

SwissProt:	O15371
IHC positive control:	Human breast carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	64 kDa
WB Positive control:	NIH/3T3 cells lysate
WB Recommended dilution:	500-3000



