

## 兔抗 TRAF1 多克隆抗体

中文名称：兔抗 TRAF1 多克隆抗体

英文名称： Anti-TRAF1 rabbit polyclonal antibody

别名： EBI6; MGC:10353

抗原： TRAF1

储存： 冷冻（-20℃）

宿主： Rabbit

相关类别： 一抗

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

**Background:**

The protein encoded by this gene is a member of the TNF receptor (TNFR) associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from various receptors of the TNFR superfamily. This protein and TRAF2 form a heterodimeric complex, which is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF2 also interacts with inhibitor-of-apoptosis proteins (IAPs), and thus mediates the anti-apoptotic signals from TNF receptors. The expression of this protein can be ind

	uced by Epstein-Barr virus (EBV). EBV infection membr ane protein 1 (LMP1) is found to interact with this an d other TRAF proteins; this interaction is thought to li nk LMP1-mediated B lymphocyte transformation to th e signal transduction from TNFR family receptors.
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	TRAF1
<b>Immunogen:</b>	Fusion protein of human TRAF1
<b>Full name:</b>	TNF receptor-associated factor 1
<b>Synonyms :</b>	EBI6; MGC:10353
<b>SwissProt:</b>	Q13077
<b>ELISA Recommended dilution:</b>	1000-2000
<b>IHC positive control:</b>	Human colon cancer and human cervical cancer
<b>IHC Recommend dilution:</b>	15-50
<b>WB Predicted band size:</b>	46 kDa
<b>WB Positive control:</b>	Human lymphoma tissue
<b>WB Recommended dilution:</b>	200-1000



